

Indirect searches for Z -like resonances at the LHC

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP08(2009)082

(<http://iopscience.iop.org/1126-6708/2009/08/082>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 59.162.23.19

The article was downloaded on 06/09/2010 at 17:17

Please note that [terms and conditions apply](#).

Indirect searches for Z' -like resonances at the LHC¹

Thomas G. Rizzo

*SLAC National Accelerator Laboratory,
2575 Sand Hill Rd., Menlo Park, CA 94025, U.S.A.*

E-mail: rizzo@slac.stanford.edu

ABSTRACT: We explore the possibility of indirectly observing the effects of Z' -like particles with electroweak strength couplings in the Drell-Yan channel at the LHC with masses above the resonance direct search reach. We find that, mostly due to statistical limitations, this is very unlikely in almost all classes of models independently of the spin of the resonance. Not unexpectedly, the one possible exception to this general result is the case of degenerate Kaluza-Klein (KK) excitations of the photon and Z that occur in some extra-dimensional models. In this special case, the strong destructive interference with the Standard Model (SM) exchanges below the resonance mass leads to a well-known significant suppression of the cross section and thus increased sensitivity to this particular new physics scenario.

KEYWORDS: Beyond Standard Model, Hadronic Colliders

ARXIV EPRINT: [0904.2534](https://arxiv.org/abs/0904.2534)

¹Work supported in part by the Department of Energy, Contract DE-AC02-76SF00515.

Contents

1	Introduction and background	1
2	Analysis	2
3	Discussion and conclusions	10

1 Introduction and background

The LHC is turning on later this year and the exploration of the TeV scale will commence soon thereafter. One of the cleanest signals for new physics that may be discovered early on is the existence of a Z' -like resonance in the Drell-Yan channel. Such resonance structures, with a wide variety of possible identities, are predicted to exist in a large number of extensions to the Standard Model(SM) [1]. Presently, for states with typical electroweak strength couplings to SM fields, current constraints from the Tevatron restrict their masses to lie above ~ 0.8 – 1 TeV [2]. Once large integrated luminosities of order ~ 100 – 300 fb $^{-1}$ are obtained at the LHC, these new resonances should be visible for masses as large as ~ 4 – 6 TeV depending on the specific details of the model. The only way to directly access states with larger masses at the LHC is to significantly increase the integrated luminosity into the few ab $^{-1}$ range as might be expected at the SLHC upgrade. Eventually, however, these direct searches will run out of steam resulting in a limited mass reach. Is there any way to see beyond this direct search reach at the LHC?

It is well-known that future e^+e^- colliders operating in the TeV energy range can indirectly probe for new Z' resonances by exploring contact interaction-like deviations from the cross sections and asymmetries predicted by the SM [3] in various final state channels. For the typical luminosities that are often discussed for these colliders, ~ 0.5 – 1 ab $^{-1}$, and making use of electron and positron beam polarizations, the masses of Z' -like states as large as ~ 10 – $20\sqrt{s}$ may be explored in this way. If these new resonance states are not too heavy it may also be possible to extract the corresponding resonance mass itself (*without* prior knowledge from the LHC) using e^+e^- data collected at several different center of mass energies [4]. Unfortunately, it seems more than likely that such colliders will not become available until at least the end of the 2020's. In the meantime we might wonder if the mass reach of the (S)LHC itself could be extended in any similar way. In this paper we will explore the possibility of indirectly observing the effects of such Z' -like states which are too massive to be seen directly as resonances at the the LHC, somewhat in analogy with what is done at e^+e^- colliders, employing data taken over a range of dilepton masses. Clearly such an analysis is only applicable to resonances with couplings to the SM fermions which are of electroweak strength or larger. We note in passing that lighter Drell-Yan resonance states with very small couplings, perhaps missed by the Tevatron, will be difficult if not impossible to observe indirectly at the LHC and so will not be a subject for discussion here.

To some extent, if we are exploring dilepton invariant masses *far* below the new resonance, one might imagine that much of this indirect search information could be extracted directly from the more conventional contact-interaction studies at the LHC [5] and this is indeed the case to some extent. However, in the more interesting (and, perhaps, more likely) case that we will consider here, the mass of the resonance is not too far above the 5σ discovery reach of the LHC so that non-interference terms as well as the finite resonance width effects also become important thus warranting a separate study. In addition to this, we note that traditional contact-interaction analyses are usually restricted to (combinations of) the familiar chiral, dimension-6, four-fermion operators introduced long ago [6] and do not allow for the possibility of other spin combinations that we must in all generality consider below.

2 Analysis

Given the rather wide selection of possible models with Drell-Yan resonances having electroweak strength couplings to the SM fermions, we will employ four sample benchmark examples in the analysis that follows: (i) a conventional Z' with electroweak couplings. Here we will employ the specific Z' of the Sequential SM (SSM) which is a standard candle for experimenters; in this case the Z' is simply the SM Z but with a significantly larger mass. Other, more realistic, Z' states with couplings of comparable magnitude will yield very similar results since this example captures the essential properties of this class of spin-1 models. (ii) A Randall-Sundrum(RS) [7] graviton with the SM fields localized to the TeV brane; to be specific we will assume that $k/M_{pl} = 0.04$ as a typical value for purposes of demonstration. Alternative values of this parameter will, of course, modify the signal rate we would obtain. This model allows us to examine the case of a spin-2 resonance which corresponds to a dimension-8 four-fermion interaction far below the (first) graviton mass. Of course the complete graviton Kaluza-Klein(KK) spectrum must be included in such an analysis; here for practical reasons we will sum over the first dozen KK states which provide the dominant contributions to the scattering amplitude in the dilepton mass range of interest at and below the first resonance peak. Similarly, (iii) we consider an R -parity violating sneutrino, $\tilde{\nu}$, which we choose for simplicity to have identical electromagnetic strength lepton number violating couplings [8] to both quarks and leptons as a typical example of a spin-0 resonance. Other choices would lead to modifications in the predicted rates below. It is important to remember that in this spin-0 case there is no interference with the usual SM exchanges. Similarly, in the KK graviton spin-2 case above, the interference with the SM exchanges also vanishes when one integrates over the full angular range. Finally, we consider (iv) the case of simultaneous, TeV-scale, Kaluza-Klein excitations of both the photon and Z with the SM fermions being 4-D [9] fields localized at the origin of the fifth dimension. In this case each of the resonances have the same couplings as do their SM counterparts except for a rescaling by a factor of $\sqrt{2}$ due to the normalization of their bulk wavefunctions. This is a very special situation due to the well-known, but somewhat unusual, destructive interference that takes place between the SM (γ, Z) and full set of KK exchange amplitudes [10] which causes a drastic reduction in the cross section with dilep-




Figure 1. 5σ resonance discovery reach for conventional Z' states at the 14 TeV LHC as a function of the integrated luminosity in the dilepton channel. The red(green, blue, magenta, cyan) curves correspond to the $Z'_{SSM}(Z'_{\psi}, Z'_{\chi}, Z'_{\eta}, Z'_{LRM})$ model cases, respectively.

ton invariant masses near ~ 0.55 that of the first KK resonances.¹ As in the RS graviton KK case, in practical calculations we will sum over the first dozen resonance states whose contributions will dominate in the dilepton mass region of interest. We note that these are only benchmark models; in some cases, e.g., that of the R-parity violating sneutrino, the overall coupling strength has been chosen to make the resonance ‘interesting’ in the mass range we are studying.

To be specific in the analysis that follows, we will only consider the LHC with $\sqrt{s} = 14$ TeV and with an integrated luminosity of 300 fb^{-1} . However, the results that we obtain can be straightforwardly translated to other effective values of \sqrt{s} and integrated luminosity. Furthermore, the CTEQ6.6 PDF’s [11] will be employed throughout the analysis below in obtaining all of our numerical results as will the usual mass-scale-dependent NLO/NNLO K-factors [12] whose typical value is $\simeq 1.25$ in the mass range of interest. The first step in our analysis is to approximately determine the usual 5σ direct discovery reach for the resonances in the benchmark set. This can be easily done by examining and extending the results from the detector collaborations themselves [13, 14]. Once these values are determined, we will consider the corresponding resonances with somewhat larger masses.

An example of this reach for conventional Z' models is shown in figure 1 as a function of the 14 TeV LHC integrated luminosity. Here we have assumed an approximate 100%

¹This is simply the result of being in the mass range above the SM Z yet below the KK mass so that the SM and KK amplitudes have nearly identical structure but opposite signs, the KKs being just heavy copies of the SM states. This destructive interference is difficult to duplicate in other model scenarios even if multiple exchanges were present since a very fine tuning of couplings would be required.

lepton ID efficiency in this and in all figures to follow. Note that a detector smearing of $\sim 1\%$, similar to that of ATLAS for electrons and CMS for muons in this mass range, has been included and, although no p_T cut is applied, the final state leptons are required to be central, i.e., with a pair rapidity $|y| \leq 2.5$ in all cases. We will employ this cut in all cases below except where as otherwise noted; these results are found to match very closely those quoted by both experimental collaborations. Here we see that in the SSM Z' model case the LHC discovery reach is ~ 5 TeV assuming a luminosity of 300 fb^{-1} . In a similar fashion, updating the estimates from our various earlier analyses, we can ascertain the 5σ discovery reaches for the other benchmark models described above. Conservatively, we find values of ~ 4 TeV for the lightest RS graviton [16], ~ 5.5 TeV [17] for the R -parity violating $\tilde{\nu}$ and ~ 5.5 – 6 TeV [18] for the first photon- Z gauge KK states, respectively.

What observables are available to search for new resonances indirectly at the LHC? There are essentially two: the dilepton invariant mass distribution, dN/dM , itself and the forward-backward asymmetry, A_{FB} , which is determined by the dilepton angular distribution. Here we focus on A_{FB} instead of the angular distribution itself since, as we will see, there are already significant statistical issues that we must face. Conventionally, the scattering angle for the $q\bar{q} \rightarrow l^+l^-$ process is defined in the center-of-mass frame (where the leptons are back-to-back) to be that between the incoming quark and the outgoing negatively charged lepton. At the LHC this quantity cannot be directly measured since we do not know from which proton the quark came [1]. However, this problem can be somewhat circumvented as quarks generally carry more momentum in the proton than do anti-quarks and thus the direction of motion, i.e., boost, of the dilepton system in the lab frame is more likely to be that of the incoming quark. Since this approach can sometimes incorrectly assign the quark direction, the resulting raw asymmetry is diluted and this must be corrected for in the analysis via Monte Carlo [20]. (We will assume that this can be done correctly and with high efficiency in all cases in our discussion below). Correspondingly, this approach requires us to place a lower bound on the rapidity of the outgoing leptons in high mass events in order to define forward and backward hemispheres to obtain A_{FB} . Here one traditionally requires [1] that the lepton pair rapidity $|y| \geq 0.8$ which removes very central events in the detector, but also results in reduced statistics thus increasing the corresponding error on the value of A_{FB} .

This is actually a more serious issue than one may initially imagine as a further problem with A_{FB} develops for the case of the very high mass dilepton pairs that we are considering: the collider kinematics leads to an upper limit on the lepton pair rapidity (in addition to the usual experimental cut of $|y| \leq 2.5$) given by the expression found in ref. [19]:

$$|y| \leq \log(\sqrt{s}/M). \quad (2.1)$$

Note that in the Drell-Yan process the fractional momenta of the initial state partons, which must satisfy $sx_1x_2 = M^2$, are given [19] by $x_{1,2} = (M/\sqrt{s})e^{\pm y}$; requiring that $0 \leq x_{1,2} \leq 1$ immediately leads to the constraint equation above. Thus, ever larger values of M for fixed \sqrt{s} force the final state leptons to be more and more central. Since here $\sqrt{s} = 14$ TeV, taking $M = 3(4, 5, 6)$ TeV as will be relevant below, leads to the constraint $|y| \leq 1.54(1.25, 1.03, 0.85)$, respectively. Since we are simultaneously requiring

that $|y| \geq 0.8$ to obtain a reasonably undiluted A_{FB} in the first place it is clear that the available phase space is very rapidly shrinking at high masses leading to rather poor available statistics. As we will see below, this implies that A_{FB} will not play a major role in indirect high mass resonance searches at the LHC even if we allow for a somewhat softer lower cut on the lepton rapidity.

Let us begin our analysis by considering the invariant mass distribution for Drell-Yan produced dileptons, dN/dM . A straightforward calculation including K-factors, efficiencies, experimental cuts and approximate detector smearing as discussed above but *neglecting* the effect of bin-to-bin $\sim \sqrt{N}$ statistical fluctuations (associated with the limited number of events in each bin and causing the apparent ‘up and down’ bin to bin changes in the plot) leads to the results shown in figure 2. Here we see several things: first, this figure verifies to some extent our previous estimates of the 5σ search reaches for the set of benchmark models that we are considering. Second, for the cases where the resonances are near or above the direct search reach, we can easily observe in some cases what appear to be deviations from the SM predictions for the dN/dM distribution in the dilepton mass range from ~ 2.5 – 4 TeV, somewhat below the resonance masses themselves. If these deviations were statistically significant (neglecting for the moment any discussion of systematic errors arising from, e.g., PDF uncertainties and electroweak corrections) by fitting the shape of this deviation to a model-dependent excitation curve we would obtain a smoking gun for the existence of a new resonance as well as a possible estimate of its mass. It is important to remember that in obtaining these results we have assumed the coupling parameters as described above. The fact that, e.g., the sneutrino is more easily visible than is the RS graviton is merely due to the choices of couplings we have made above for the assumed values of the resonance masses under study for the different models. What we have shown by this analysis is that, in the absence of statistical fluctuations, some resonances will produce visible deviations in the dilepton mass distribution from the SM expectations in the region below the resonance mass. As we will now show, these fluctuations lead to a ‘washing-out’ of this effect rendering indirect observation essentially impossible.

In reality, however, the true experimental picture will be much more similar to the corresponding results shown in figure 3 where the statistical fluctuations are now explicitly included. Here we can still observe (at the 5σ level in comparison to the SM background) resonances themselves (at the corresponding mass reach values quoted above) but the statistical fluctuations in the data below the resonance peak now make all of the distributions look rather similar. A careful analysis does show some sensitivity to the higher mass resonances (which are quite close to being observable directly) in the case of KK gauge excitations as these have a very special associated signature due to the strong destructive interference in this mass range as discussed above [10]. Even with the statistical fluctuations present it is possible to observe that the prediction for the gauge KK case lies systematically *below* that of the SM. One could argue that this bad situation with respect to the statistical fluctuations masking any significant effect might be improved by increasing the width of the dilepton mass bins to increase statistics and decrease the effects of fluctuations. In this case, however, we only gain in statistics at the price of reduced sensitivity as can be seen by the example shown in figure 4, where the bin sizes are now




Figure 2. Top panel: idealized dilepton invariant mass distributions, neglecting the effects of statistical fluctuations (i.e., finite statistics accounting only for the overall rate and not the $\sim \sqrt{N}$ uncertainty effects in any given bin), for the SM (black), the SSM with a 5 TeV Z' (red), a photon/ Z gauge KK state with a mass of 5 TeV (cyan), an RS graviton with a 4 TeV mass (blue) and a 4 TeV R -parity violating sneutrino (green). Bottom panel: same as above but now with all the resonance masses increased by 1 TeV. A detector smearing of $\sim 1\%$, similar to that of ATLAS for electrons and CMS for muons in this mass range, has been included and the final state leptons are required to be central, i.e., $|y| \leq 2.5$ in all cases as discussed in the text. Unless otherwise stated our bin width is 1% of the central mass value for each bin and NLO/NNLO K-factors are included.




Figure 3. Same as the previous figure but now showing the effect of the bin-to-bin $\sim \sqrt{N}$ statistical fluctuations in the number of events due to finite luminosity.

increased by a relative factor of 5, with similar results found to hold in other rescaling cases. It thus seems quite unlikely that new resonances can be indirectly observed at the LHC except in exceptional circumstances where they have large effects on data in the dilepton mass range far below the resonance mass itself.




Figure 4. Same as the lower panel in previous figure but now showing the effect of increasing the width of all the bins by a factor of 5 to increase statistics.

An alternative approach to using the dN/dM distribution itself is to use the same data to form the associated cumulative distribution which displays the total number of events above some minimum dilepton mass cutoff. Such distributions certainly have added statistical power as well as increased sensitivity to the high mass region of interest to us and are not sensitive to bin-size effects. Examples of these types of distributions using our benchmark models for both sets of assumed masses are shown in figure 5. In the top panel, all of the benchmark models are clearly seen above the SM background as would be expected from our earlier discussion. In the lower panel, the 5 TeV sneutrino and the 6 TeV gauge KK benchmarks may possibly lie significantly above the SM background while the other benchmarks appear to lie much lower. For comparison purposes the case of a 6 TeV sneutrino is also shown and the corresponding histogram is seen to lie quite close to the SM background expectations. In order to make a direct comparison of the benchmark models, consider, for example, the cumulative number of expected events we find above a dilepton mass of 4 TeV (subject to very large statistical fluctuations as well as possibly significant systematic uncertainties) is 1.0(1.8, 1.4, 3.9, 7.2, 1.7) for the SM(6 TeV SSM, 5 TeV RS graviton, 6 TeV gauge KK, 5 TeV sneutrino and 6 TeV sneutrino) cases, respectively, none of which remotely rises close to the level of a significant signal [21] except possibly for the case of the 5 TeV sneutrino. Clearly the cumulative distributions do not show any obvious evidence for sensitivity to new resonances beyond the direct LHC discovery reach even when potentially important systematic effects are neglected. This implies that except in exceptional circumstances new resonances cannot be indirectly detected using this method.




Figure 5. Same as in figure 3 but now showing the *cumulative* number of events above a minimum value of the dilepton invariant mass. In the lower panel an additional (lower) green histogram is also present for the R -parity violating sneutrino case assuming a resonance mass of 6 TeV. Note that statistical fluctuations are included here in both panels.

Let us now return to the issue of the forward-backward asymmetry, A_{FB} , discussed above. In principle, this quantity shows a reasonable sensitivity to heavy resonances but is limited in applicability by reduced statistics. This can be most easily seen by examining the results presented in the top panel of figure 6 which shows (without errors) the expected

values of A_{FB} in the dilepton mass range of interest for the SM as well as for the SSM and gauge KK models for several different values of the resonance masses. Clearly A_{FB} deviates significantly from the SM expectations in the region below the mass of the new resonance in all cases. However, as we will see, the statistical power of A_{FB} in this same mass range is quite poor. To demonstrate the importance of a cut of $|y| \geq 0.8(0.5)$ on the available statistics we note that in the lowest dilepton mass bin considered here, near ~ 1 TeV, we find 248 events in the SM case before applying this rapidity cut but only 105(155) events afterwards. In this mass bin, these cuts yield the asymmetry values of $A_{FB} = 0.388 \pm 0.090(0.346 \pm 0.075)$ which show that the (statistical only) errors are already quite large. Above this dilepton mass the severity of the lower rapidity cut becomes far more significant and the size of the error on the asymmetry grows quite rapidly.² This result is explicitly seen in the lower panel in figure 6 for either choice of the value of the lower cut on $|y|$. From this figure, however, we can easily imagine that if the mass of the new resonance were significantly less, 1.5–2 TeV say, it would be possible that A_{FB} measurements below the peak would provide significant sensitivity to new physics. In fact, it is well-known that for the new gauge boson case such data provides important information on the couplings of Z' -like states [1].

By combining the A_{FB} information for all dilepton masses above ~ 1 TeV we can determine the added sensitivity to new resonance states beyond the direct discovery search reach. Clearly such sensitivity will be dependent upon the specific nature of this new resonance and the precise lower bound on the rapidity applied to the final state leptons. However, scanning over the various benchmark models and possible resonance masses we find that the total contribution to the χ^2 from the deviation in the value of A_{FB} to be always less than ~ 0.3 – 0.5σ and generally far smaller. This is not too surprising given the results shown in figure 6. Thus it is quite clear that the additional information obtained from the dilepton angular distribution will not be very helpful in indirect searches for new resonances in the dilepton mass range of interest to us here above 4–5 TeV. Thus we conclude that such searches must essentially rely on the dilepton mass distribution except in very special circumstances.

3 Discussion and conclusions

In this paper we have explored the possibility of indirectly discovering resonance states in the Drell-Yan channel at the LHC for particles of various spins when their masses are just beyond the direct discovery reach. Generally, we find this task to be quite difficult if not impossible even in the limit where systematic effects, such as PDF uncertainties, are neglected. The only exception to this result is the rather unusual circumstance of nearly degenerate photon and Z Kaluza-Klein excitations. In particular, in the course of this analysis, we found that the forward-backward asymmetry, A_{FB} , of the dilepton pair was generally not a useful observable for performing such searches due to the large associated

²It is important to remember that the severity of the rapidity cut increases with the dilepton mass due to the kinematic upper limit given in eq. (2.1).




Figure 6. The upper panel shows the predicted values of A_{FB} in the SM (yellow histogram), for the SSM with a Z' mass of 4(5, 6) TeV corresponding to the red (green, blue) histograms, and for the case of gauge KK excitations with mass of 5(6, 7) TeV, corresponding to the magenta (cyan, solid) histograms, respectively. For purposes of demonstration, a cut of $|y| \geq 0.5$ has been applied in all cases. The lower panel shows the statistical error on the SM value of A_{FB} as a function of the dilepton mass assuming a cut of $|y| \geq 0.5$ (0.8) corresponding to the red (blue) histogram. An integrated luminosity of 300 fb^{-1} has been assumed in the lower panel.

statistical errors in the mass range of interest above ~ 2 – 2.5 TeV. Essentially, these indirect searches must rely solely on the dilepton mass distribution. Except for possibly the special gauge KK case, where significant modifications are observed in the dilepton invariant mass distribution at masses far below the peak due to strong interference with SM exchanges, new resonances are found to only perturb the mass distribution significantly in the neighborhood just below the peak itself where the statistics are commonly quite poor.

Hopefully new Drell-Yan resonances will be relatively light and will soon be discovered at the LHC.

Acknowledgments

The author would like to thank J. Wells for raising the important issue of indirect resonance searches at the LHC and for early discussions on the work presented here. The author would also like to thank J.L. Hewett for discussions and the CERN Theory Group, where part of this work was done, for its hospitality.

References

- [1] For recent reviews and original references, see P. Langacker, *The Physics of Heavy Z' Gauge Bosons*, [arXiv:0801.1345](#) [SPIRES];
 T.G. Rizzo, *Z' phenomenology and the LHC*, [hep-ph/0610104](#) [SPIRES];
 M.S. Carena, A. Daleo, B.A. Dobrescu and T.M.P. Tait, *Z' gauge bosons at the Fermilab Tevatron*, *Phys. Rev. D* **70** (2004) 093009 [[hep-ph/0408098](#)] [SPIRES];
 M. Dittmar, A.-S. Nicollerat and A. Djouadi, *Z' studies at the LHC: an update*, *Phys. Lett. B* **583** (2004) 111 [[hep-ph/0307020](#)] [SPIRES];
 for some classic reviews of new gauge boson physics, see A. Leike, *The Phenomenology of extra neutral gauge bosons*, *Phys. Rept.* **317** (1999) 143 [[hep-ph/9805494](#)] [SPIRES];
 J.L. Hewett and T.G. Rizzo, *Low-Energy Phenomenology of Superstring Inspired E_6 Models*, *Phys. Rept.* **183** (1989) 193 [SPIRES];
 M. Cvetič and S. Godfrey, *Discovery and identification of extra gauge bosons*, [hep-ph/9504216](#) [SPIRES];
 T.G. Rizzo, *Extended gauge sectors at future colliders: report of the new gauge boson subgroup*, in *Proceedings of the 1996 DPF/DPB Summer Study on New Directions in High-energy Physics*, Snowmass, Colorado, U.S.A., June 25 – July 12 1996, pg. NEW136, [hep-ph/9612440](#) [SPIRES].
- [2] For recent constraints from the Tevatron, see CDF collaboration, T. Aaltonen et al., *A search for high-mass resonances decaying to dimuons at CDF*, *Phys. Rev. Lett.* **102** (2009) 091805 [[arXiv:0811.0053](#)] [SPIRES].
- [3] S. Riemann, *Prospects to detect a Z' with the LC*, [hep-ph/9710564](#) [SPIRES];
 S. Godfrey, *Comparison of discovery limits for extra Z bosons at future colliders*, *Phys. Rev. D* **51** (1995) 1402 [[hep-ph/9411237](#)] [SPIRES];
 T.G. Rizzo, *Contact interaction searches at the linear collider: energy, luminosity and positron polarization dependencies*, [hep-ph/0303056](#) [SPIRES].
- [4] T.G. Rizzo, *An exploration of below threshold Z' mass and coupling determinations at the NLC*, *Phys. Rev. D* **55** (1997) 5483 [[hep-ph/9612304](#)] [SPIRES].

- [5] For a recent overview, see for example P. Spagnolo, *Prospects for discovering new gauge bosons, extra dimensions and contact interaction at the LHC*, [arXiv:0810.0414](#) [SPIRES]; for details of recent studies by ATLAS and CMS, see refs. [13] and [14] below.
- [6] E. Eichten, K.D. Lane and M.E. Peskin, *New Tests for Quark and Lepton Substructure*, *Phys. Rev. Lett.* **50** (1983) 811 [SPIRES].
- [7] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)] [SPIRES].
- [8] H.K. Dreiner, P. Richardson and M.H. Seymour, *Resonant slepton production*, [hep-ph/9903419](#) [SPIRES];
 J. Kalinowski, R. Ruckl, H. Spiesberger and P.M. Zerwas, *Supersymmetry with R-parity breaking: contact interactions and resonance formation in leptonic processes at LEP2*, *Phys. Lett. B* **406** (1997) 314 [[hep-ph/9703436](#)] [SPIRES]; *R-parity violating SUSY signals in lepton pair production at the Tevatron*, *Phys. Lett. B* **414** (1997) 297 [[hep-ph/9708272](#)] [SPIRES];
 J. Erler, J.L. Feng and N. Polonsky, *A wide scalar neutrino resonance and $b\bar{b}$ production at LEP*, *Phys. Rev. Lett.* **78** (1997) 3063 [[hep-ph/9612397](#)] [SPIRES];
 B.C. Allanach, H.K. Dreiner, P. Morawitz and M.D. Williams, *Single sneutrino/slepton production at LEP2 and the NLC*, *Phys. Lett. B* **420** (1998) 307 [[hep-ph/9708495](#)] [SPIRES];
 S. Bar-Shalom, G. Eilam and A. Soni, *R-parity violation and CP-violating and CP-conserving spin asymmetries in $\ell^+\ell^- \rightarrow \tilde{\nu} \rightarrow \tau^+\tau^-$: probing sneutrino mixing at $\ell^+\ell^-$* , *Phys. Rev. Lett.* **80** (1998) 4629 [[hep-ph/9802251](#)] [SPIRES]; *R-parity violation, sneutrino mixing phenomena and CP- violation in $p\bar{p} \rightarrow \tilde{\nu} \rightarrow \ell^+\ell^- + X$* , *Phys. Rev. D* **59** (1999) 055012 [[hep-ph/9804339](#)] [SPIRES];
 J.L. Feng, J.F. Gunion and T. Han, *R-parity violation and sneutrino resonances at muon colliders*, *Phys. Rev. D* **58** (1998) 071701 [[hep-ph/9711414](#)] [SPIRES].
- [9] I. Antoniadis, K. Benakli and M. Quirós, *Production of Kaluza-Klein states at future colliders*, *Phys. Lett. B* **331** (1994) 313 [[hep-ph/9403290](#)] [SPIRES];
 P. Nath, Y. Yamada and M. Yamaguchi, *Probing the nature of compactification with Kaluza-Klein excitations at the Large Hadron Collider*, *Phys. Lett. B* **466** (1999) 100 [[hep-ph/9905415](#)] [SPIRES];
 K.-m. Cheung and G.L. Landsberg, *Kaluza-Klein states of the standard model gauge bosons: constraints from high energy experiments*, *Phys. Rev. D* **65** (2002) 076003 [[hep-ph/0110346](#)] [SPIRES].
- [10] For a detailed study, see G. Azuelos and G. Polesello, *Prospects for the detection of Kaluza-Klein excitations of gauge bosons in the ATLAS detector at the LHC*, *Eur. Phys. J. C* **39S2** (2005) 1 [SPIRES].
- [11] P.M. Nadolsky et al., *Implications of CTEQ global analysis for collider observables*, *Phys. Rev. D* **78** (2008) 013004 [[arXiv:0802.0007](#)] [SPIRES].
- [12] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $O(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017 [[hep-ph/0609070](#)] [SPIRES];
 see, for example, F. Petriello and S. Quackenbush, *Measuring Z' couplings at the CERN LHC*, *Phys. Rev. D* **77** (2008) 115004 [[arXiv:0801.4389](#)] [SPIRES];
 P. Mathews and V. Ravindran, *NLO QCD corrections to Drell-Yan in TeV-scale gravity models*, *Nucl. Phys. Proc. Suppl.* **157** (2006) 251 [SPIRES].

- [13] THE ATLAS collaboration, G. Aad et al., *Expected Performance of the ATLAS Experiment — Detector, Trigger and Physics*, [arXiv:0901.0512](https://arxiv.org/abs/0901.0512) [SPIRES].
- [14] *CMS Physics Technical Design Report*, <https://cmsdoc.cern.ch/cms/cpt/tdr/>;
CMS collaboration, G.L. Bayatian et al., *CMS technical design report, volume II: physics performance*, *J. Phys. G* **34** (2007) 995 [SPIRES].
- [15] P. Mathews and V. Ravindran, *NLO QCD corrections to Drell-Yan in TeV-scale gravity models*, *Nucl. Phys. Proc. Suppl.* **157** (2006) 251 [SPIRES].
- [16] H. Davoudiasl, J.L. Hewett and T.G. Rizzo, *Experimental probes of localized gravity: on and off the wall*, *Phys. Rev. D* **63** (2001) 075004 [[hep-ph/0006041](https://arxiv.org/abs/hep-ph/0006041)] [SPIRES]; *Phenomenology of the Randall-Sundrum Gauge Hierarchy Model*, *Phys. Rev. Lett.* **84** (2000) 2080 [[hep-ph/9909255](https://arxiv.org/abs/hep-ph/9909255)] [SPIRES].
- [17] J.L. Hewett and T.G. Rizzo, *Resonant slepton production at hadron colliders in R- parity violating models*, [hep-ph/9809525](https://arxiv.org/abs/hep-ph/9809525) [SPIRES].
- [18] T.G. Rizzo, *Testing the nature of Kaluza-Klein excitations at future lepton colliders*, *Phys. Rev. D* **61** (2000) 055005 [[hep-ph/9909232](https://arxiv.org/abs/hep-ph/9909232)] [SPIRES].
- [19] For a review of the dilepton production kinematics at hadron colliders and the origin of the following expression, see V.D. Barger and R.J.N. Phillips, *Collider physics*, Frontiers in Physics **71**, Addison-Wesley, Redwood City, U.S.A. (1987);
and also E. Eichten, I. Hinchliffe, K.D. Lane and C. Quigg, *Super Collider Physics*, *Rev. Mod. Phys.* **56** (1984) 579 [*Addendum ibid.* **58** (1986) 1065] [SPIRES].
- [20] For a nice discussion, see R. Cousins, J. Mumford, J. Tucker and V. Valuev, *Measurement of Forward-Backward Asymmetry of Simulated and Reconstructed $Z' \rightarrow \mu^+ \mu^-$ Events in CMS*, CMS-NOTE-2005-022.
- [21] G.J. Feldman and R.D. Cousins, *A Unified Approach to the Classical Statistical Analysis of Small Signals*, *Phys. Rev. D* **57** (1998) 3873 [[physics/9711021](https://arxiv.org/abs/physics/9711021)] [SPIRES].